arXiv:1212.1426v1 [math.AG] 6 Dec 2012

ON BÜCHI'S K3 SURFACE

MICHELA ARTEBANI, ANTONIO LAFACE, AND DAMIANO TESTA

Abstract. We study the geometry of Büchi's K3 surface showing that the rational points of this surface are Zariski-dense.

Introduction

The Büchi problem is a well known question in number theory: it asks whether there exists a positive integer n such that any sequence $\{x_i^2\}$ of n integer squares with

$$(x_{i+2}^2 - x_{i+1}^2) - (x_{i+1}^2 - x_i^2) = 2 \qquad \text{for } i \in \{1, \dots, n-2\} \tag{0.1}$$

is necessarily a sequence of squares of the form $\{(x+i)^2\}$, with $i \in \{1, \dots, n\}$ and x an integer. A positive answer to Büchi's problem would imply, using the negative answer to Hilbert's Tenth Problem by Yu. Matiyasevich, that there is no algorithm to decide whether a system of diagonal quadratic forms with integer coefficients represents an integer tuple (see [PPV10]).

A solution of (0.1) is *trivial* if it is given by a sequence of squares of consecutive integers, and *non-trivial* otherwise. It is known that if $n \leq 4$ there are non-trivial integral solutions of (0.1). In this paper we prove that the set of non-trivial *rational* solutions, when $n = 5$, is infinite and Zariski-dense in the corresponding algebraic surface. More specifically, since we are interested in the rational points of (0.1), we consider the projective surface X of $\mathbb{P}^5$ defined by the homogeneous equations:

$$X\colon \quad \begin{cases} x_3^2 - 2x_2^2 + x_1^2 &= \quad 2x_0^2 \\ x_4^2 - 2x_3^2 + x_2^2 &= \quad 2x_0^2 \\ x_5^2 - 2x_4^2 + x_3^2 &= \quad 2x_0^2. \end{cases} \tag{0.2}$$

As we shall see, the surface X is the Kummer surface associated to an abelian surface which is isogenous, over the complex numbers, to the cartesian product $E \times E$, where E is an elliptic curve without complex multiplication. By means of this fact we compute an explicit basis for the Picard group of X and construct one-parameter families of non-trivial rational solutions of the Büchi problem for $n = 5$. Using the Mukai correspondence we prove that the surface X is an irreducible component of the moduli space of stable vector bundles on X itself: the modular point of view allows us to interpret rational points on X in terms of rank two vector bundles and opens the possibility of using techniques from derived category

---

2000 *Mathematics Subject Classification.* 14J28, 11D41.

Key words and phrases. Büchi problem, Kummer surfaces, rational points.

The first author has been partially supported by Proyecto FONDECYT Regular N. 1110249. The second author has been partially supported by Proyecto FONDECYT Regular N. 1110096. The third author has been partially supported by EPSRC grant EP/F060661/1.

and moduli spaces to attack the Büchi problem. It also raises the question of which vector bundles correspond to integral solutions of the system (0.2) and the related question of finding a description in terms of vector bundles for the points lying on the curve with equation $x_0 = 0$ on X.

The paper is organised as follows. In Section 1 we introduce Büchi surfaces and their linear automorphisms groups. In Section 2 we focus on the Büchi K3 surface X showing that it is a Kummer surface: the associated abelian variety is isogenous to a cartesian product $E \times E$, where E is an elliptic curve without complex multiplication. As a consequence we determine the lattice structure of the Picard group of X. Section 3 is devoted to the study of two elliptic pencils on X defined over the rational numbers. This allows us to produce one-parameter solutions of (0.2) and to show that the rational points of X are Zariski-dense. We determine in Section 4 an interpretation of the surface X as an irreducible component of the moduli space of stable vector bundles on X itself. Finally, Section 5 analyzes the Galois representation of the elliptic curve E and its set of supersingular primes: we use these results to compute the local Zeta function of X at such primes.

Acknowledgements. The authors would like to thank Kieran O'Grady and Daniel Huybrechts for their suggestions and their help with the Mukai correspondence and moduli spaces of sheaves on K3 surfaces.

1. Büchi surfaces

Let n be an integer with $n \geq 3$. The Büchi surface X_n is the zero locus in $\mathbb{P}^n$ of the $n-2$ quadrics of equations

$$x_{i+2}^2 - 2\,x_{i+1}^2 + x_i^2 - 2\,x_0^2 = 0 \qquad \text{for } i \in \{1, \dots, n-2\}. \tag{1.1}$$

Observe that X_3 is a quadric, X_4 is a del Pezzo surface of degree four, X_5 is a K3 surface and X_n, $n \geq 6$ is of general type. A first attempt to find rational points on X_n can be done by producing a parametrization of the surface when $n \leq 4$. The del Pezzo surface X_4 can be parametrized by the system of plane cubics passing through the five points

$$(1:0:0) \quad (0:1:0) \quad (0:0:1) \quad (3:-3:2) \quad (3:-1:1).$$

Explicitly we have the following parametrization of X_4:

$$\begin{aligned}
x_0 &= -2a^2b - 6ab^2 - 3a^2c + 9b^2c + 9ac^2 + 9bc^2\\
x_1 &= -12ab^2 + 3a^2c - 18abc + 9b^2c - 18ac^2\\
x_2 &= 2a^2b - 6ab^2 - 24abc - 9ac^2 + 9bc^2\\
x_3 &= -4a^2b - 3a^2c + 6abc - 9b^2c - 18bc^2\\
x_4 &= -6a^2b - 6ab^2 - 6a^2c - 18b^2c + 9ac^2 + 27bc^2
\end{aligned}$$

which immediately proves the existence of infinitely many non-trivial rational solutions of the Büchi's problem for four variables. Indeed, not only the set of rational points of X_4 is infinite, but even the set of points that are integral with respect to the divisor $x_0 = 0$ is infinite (see [PPV10]). The surface X_4 contains 16 lines whose parametric equations in the affine chart $x_0 \neq 0$ are:

$$x_i = \pm(t + i), \quad i = 1, 2, 3, 4.$$

The surface X_5 is the double cover of X_4 branched along the smooth genus 5 curve $B \in |-2K_{X_4}|$ defined by $2x_4^2 - x_3^2 + 2x_0^2 = 0$ (the covering map is the projection on the first four coordinates). Since the lines of X_4 are tangent to the branch locus B, the inverse image of each line in the double cover X_5 decomposes in the union of two lines. Thus X_5 contains 32 lines, whose parametric equations are:

$$x_i = \pm(t+i), \quad i = 1, 2, 3, 4, 5. \tag{1.2}$$

Observe that these are the only lines in X_5, since the covering map is linear, and that they contain the trivial solutions of Büchi problem.

1.1. The linear automorphism group of X_n.

A *scalar permutation* is an automorphism of the projective space $\mathbb{P}^r$ whose matrix with respect to the natural homogeneous coordinates only has one non-zero entry in each row and in each column. Equivalently, a scalar permutation is an automorphism of projective space fixing the set of coordinate points (namely, those points of projective space with a unique non-zero coordinate).

Let $\tau\colon \mathbb{P}^n \to \mathbb{P}^n$ be the involution defined by

$$\tau(x_0 : \cdots : x_n) = (x_0 : x_n : x_{n-1} : \cdots : x_2 : x_1);$$

the automorphism τ induces an involution of X_n that we denote by the same symbol.

Lemma 1.1. *Assume that n is an integer with $n \geq 4$ and that the characteristic of the ground field is different from two and three. The linear automorphism group of X_n consists of scalar permutations.*

Proof. The surface X_n is defined by diagonal quadrics. If $\mathcal{Q} \subset \mathbb{P}\big(I(X_n)_2\big)$ is a set of quadrics vanishing on X_n, denote by $\Sigma_{\mathcal{Q}}$ the intersection of the singular loci of the quadrics in $\mathcal{Q}$, and note that $\Sigma_{\mathcal{Q}}$ is a coordinate linear subspace. Therefore, the linear automorphism group of X_n permutes the set of linear subspaces $\big\{\Sigma_{\mathcal{Q}} \mid \mathcal{Q} \subset \mathbb{P}\big(I(X_n)_2\big)\big\}$. Hence, to prove the assertion, it suffices to show that all coordinate points of $\mathbb{P}^n$ appear among the linear subspaces $\Sigma_{\mathcal{Q}}$. For each $i \in \{0, \ldots, n\}$ let p_i be the point of $\mathbb{P}^n$ whose only non-zero coordinate is x_i; we shall determine a set $\mathcal{Q}_i$ of quadrics vanishing on X_n such that $\Sigma_{\mathcal{Q}_i} = \{p_i\}$. Using the involution τ, it suffices to determine $\mathcal{Q}_i$ for those indices i satisfying $i \leq n+1-i$.

For the coordinate point p_0 we can choose $\mathcal{Q}_0$ to consist of all the quadrics whose equations are the pairwise differences of the equations in (1.1).

For $i \in \{1, \ldots, n\}$ we denote by $\mathcal{Q}'_i$ the set of all the quadrics whose equations are those among the equations in (1.1) not involving the monomial x_i^2.

The choice $\mathcal{Q}_i = \mathcal{Q}'_i$ satisfies the requirements for the indices $i \leq \frac{n+1}{2}$ in the set $\{1, 4, 5, \ldots, n-4, n-3, n\}$.

For the coordinate point p_2 we can choose $\mathcal{Q}_2$ to consist of $\mathcal{Q}'_2$ together with the quadric defined by $x_1^2 - 3x_3^2 + 2x_4^2 = 2x_0^2$.

For the coordinate point p_3 we can choose $\mathcal{Q}_3$ to consist of $\mathcal{Q}'_3$ together with the quadric defined by $2x_1^2 - 3x_2^2 + x_4^2 = 6x_0^2$. □

Theorem 1.2. *If n is an integer with $n \geq 3$ and if the characteristic of the ground field is at least $2n+1$, then the group of linear automorphisms of X_n is generated by the sign changes of the variables and the involution τ.*

Proof. By Lemma 1.1 it suffices to show that the only permutations of the coordinates fixing X_n are those determined by τ. Let $\overline{X}_n \subset \mathbb{P}^n$ be the linear subspace with equations

$$\begin{cases} x_1 - 2x_2 + x_3 & = & 2x_0 \\ \qquad x_2 - 2x_3 + x_4 & = & 2x_0 \\ \qquad\qquad \ddots & \vdots & \\ \qquad\qquad x_{n-2} - 2x_{n-1} + x_n & = & 2x_0 \end{cases}$$

and observe that the result follows if we show that the only automorphisms of $\overline{X}_n$ induced by permutations of the coordinates are the identity and τ. It is immediate to check that the only the point lying on $\overline{X}_n$ and obtained from $b_0 = (0 : 1 : 1 : \cdots : 1)$ by permuting its coordinates is the point b_0 itself. Thus, the permutations of the coordinates acting on $\overline{X}_n$ must fix the first coordinate x_0. Let i be an index in $\{1, \ldots, n\}$ and denote by b_i the unique point in the intersection of $\overline{X}_n$ with the linear space with equations $x_0 = x_i = 0$. The coordinates of the point b_i are $(0 : i-1 : i-2 : \cdots : i-n)$. By construction, the permutations of the coordinates inducing automorphisms of $\overline{X}_n$ fix the set of points $\mathcal{B} = \{b_1, b_2, \ldots, b_n\}$. Let σ be an automorphism of $\overline{X}_n$ induced by a permutation of the coordinates. The point $\sigma(b_1)$ lies in $\mathcal{B}$; by the assumption that the characteristic of the ground field is at least $2n+1$, it follows that the n non-zero coordinates and their n negatives are all distinct. Since the only two points in $\mathcal{B}$ with this property are b_1 and b_n, we obtain that $\sigma(b_1) \in \{b_1, b_n\}$. Composing σ with τ if necessary, we reduce to the case in which $\sigma(b_1) = b_1$. Again, since all the non-zero coordinates of b_1 are distinct, we conclude that σ is the identity and the result is proved. □

Remark 1.3. Each surface X_n is a complete intersection of quadrics. By the adjunction formula we see that the canonical divisor X_n is linearly equivalent to $(n-5)H$, where H is the hyperplane section of X_n. Every automorphism of X_n must preserve the (anti-)canonical linear series. Moreover, when $n \neq 5$, the canonical divisor is a non-zero multiple of the complete linear series that embeds X_n in $\mathbb{P}^n$. Since the Picard group of X_n is torsion-free, it follows that the automorphism group of X_n consists just of the linear automorphisms of X_n, in the case $n \neq 5$.

If $n = 5$ the previous argument is no longer true, since K_{X_5} is trivial. In this case the automorphism group of X_5 has infinite order and we will show in Remark 3.2 that this is true even for the subgroup $\mathrm{Aut}(X_5)_{\mathbb{Q}}$ of automorphisms defined over the rationals.

2. The Kummer surface

From now on we denote by X the Büchi K3 surface X_5.

2.1. The 32 lines.

The surface X contains four reducible hyperplane sections of equations:

$$x_2 \pm x_1 \pm x_0 = 0,$$

one for each choice of sign. Each such equation cuts 8 lines on X and all the 32 lines of X are contained in these hyperplanes. By using equations (1.2) one sees that the intersection graph of these lines on X is bipartite of type $(16, 16)$.

Proposition 2.1. *The surface X is the Kummer surface of the Jacobian surface JC, where C is the genus two curve of equation*

$$y^2 = (x-2)\,(x-1)\,x\,(x+1)\,(x+2).$$

Proof. Since X contains 16 disjoint rational curves, then it is a Kummer surface by [Nik75, Theorem 1]. Each line L on X intersects exactly six other disjoint lines $L_1, \dots, L_6$. The morphism associated to the divisor $2L + \sum_{i=1}^{6} L_i$ maps X onto a quartic surface in $\mathbb{P}^3$ with 16 nodes which are images of 16 disjoint lines of X containing the six L_i's. The hyperplane section corresponding to the previous divisor is a trope of the quartic surface, that is it intersects the surface along a double conic containing six nodes. The double cover of such conic branched along the six nodes is a curve C genus two such that $X = \mathrm{Km}(JC)$ (see [Bea96, Exercise VIII.22, (5)]). Thus C is isomorphic to the double cover of L branched along the points $L \cap L_i$. An explicit computation gives the above equation for C. $\square$

Remark 2.2. By looking at the right hand polynomial of the previous equation for C, it is clear that C is isomorphic to any genus 2 curve defined by an equation $y^2 = P(x)$, such that the five roots of $P(x)$ are in arithmetic progression.

Lemma 2.3. *The automorphism group of C is isomorphic to the order 8 dihedral group D_4 and it is generated by the following automorphisms:*

$$\sigma_1(x,y) = (-x, iy), \qquad \sigma_2(x,y) = \left(\frac{2}{x}, \frac{2\sqrt{2}y}{x^3}\right).$$

Proof. The full automorphism group of C over the complex numbers can be computed by means of Magma [BCP97]. The program calculates the Cardona-Quer-Nart-Pujola invariants of a genus two curve and makes use of the classification of the automorphism groups in terms of such invariants. The group $\mathrm{Aut}(C)$ is isomorphic to D_4 and σ_1, σ_2 are clearly two generators. $\square$

2.2. The associated abelian surface.

Let k be a field of characteristic different from two. Let F be an elliptic curve over a field k with equation $F\colon y^2 = f(x)$; we denote by F_d the quadratic twist of F by $d \in k^*$ defined by $F\colon dy^2 = f(x)$.

Let E and E' be the elliptic curves over the rational numbers with equations:

$$E\colon y^2 = x^3 - 35\,x^2 + 35\,x - 1 \qquad E'\colon y^2 = x^3 + \frac{5}{9}\,x^2 - \frac{5}{9}\,x - 1.$$

The elliptic curves E and E' have j-invariants $\frac{2744000}{9}$ and $-\frac{8000}{81}$, respectively; they are isogenous under the degree two isogeny defined by the rational two-torsion point of E (or equivalently of E').

The automorphism group of C contains two distinct conjugacy classes of non-hyperelliptic involutions, with representatives σ_2 defined over $\mathbb{Q}(\sqrt{2})$ and $\sigma_1 \circ \sigma_2$ defined over $\mathbb{Q}(\sqrt{-2})$. Observe that the involution $\sigma_1^2 \circ \sigma_2$ is obtained applying the non-trivial automorphism in $\mathrm{Gal}(\mathbb{Q}(\sqrt{2})/\mathbb{Q})$ to the coefficients of the involution σ_2. Denote by φ and $\overline{\varphi}$ the quotients

$$\varphi\colon C \to C/\langle\sigma_2\rangle \qquad \text{and} \qquad \overline{\varphi}\colon C \to C/\langle\sigma_1^2 \circ \sigma_2\rangle.$$

The curves $C/\langle\sigma_2\rangle$ and $C/\langle\sigma_1^2 \circ \sigma_2\rangle$ have genus one.

In the following proposition, we use the Weil restriction of a curve. In general, let L/K be a finite field extension; the Weil restriction functor WR is a functor from schemes X defined over L to schemes defined over K, having the property that L-rational points of X are in functorial bijection with K-rational points on $\mathrm{WR}(X)$. We simply use the initial properties of Weil restriction, and we refer the interested reader to the book [BLR90, Chapter 7.6] for more details. In our case, it is easy to compute explicit equations for (an affine open subset of) the Weil restriction $\mathrm{WR}(E_{\sqrt{2}})$ from $L = \mathbb{Q}(\sqrt{2})$ to $K = \mathbb{Q}$ of the elliptic curve $E_{\sqrt{2}}$: substituting $x_1 + \sqrt{2}x_2$ and $y_1 + \sqrt{2}y_2$ for x and y in the equation of $E_{\sqrt{2}}$ and equating coefficients of powers of $\sqrt{2}$ we find the equations

$$\mathrm{WR}(E_{\sqrt{2}})\colon \quad \left\{ \begin{array}{rcl} 4y_1y_2 & = & x_1^3 + 6x_1x_2^2 - 35x_1^2 - 70x_2^2 + 35x_1 - 1 \\ y_1^2 + 2y_2^2 & = & 3x_1^2x_2 + 2x_2^3 - 70x_1x_2 + 35x_2 \end{array} \right.$$

for an affine open subvariety of $\mathrm{WR}(E_{\sqrt{2}})$.

Proposition 2.4. *There are isomorphisms*

$$E_{\sqrt{2}} \simeq C/\langle \sigma_2 \rangle \qquad \text{and} \qquad E_{-\sqrt{2}} \simeq C/\langle \sigma_1^2 \circ \sigma_2 \rangle,$$

so that the Jacobian surface JC is isogenous to the abelian surface $\mathrm{WR}(E_{\sqrt{2}})$. A pair of dual degree four isogenies is determined by

$$\left\{ \begin{array}{rccc} \hat{\psi}\colon & JC & \longrightarrow & \mathrm{WR}(E_{\sqrt{2}}) \\ & \omega & \longmapsto & (\varphi, \overline{\varphi})_*(\omega) \end{array} \right. \quad \text{and} \quad \left\{ \begin{array}{rccc} \psi'\colon & \mathrm{WR}(E_{\sqrt{2}}) & \longrightarrow & JC \\ & (p,q) & \longmapsto & (\varphi, \overline{\varphi})^*(p,q). \end{array} \right.$$

Proof. In coordinates, the map φ is the rational map

$$(x,y) \longmapsto \left(\left(\frac{x-\sqrt{2}}{x+\sqrt{2}} \right)^2, \frac{8\sqrt{2}y}{(x+\sqrt{2})^3} \right)$$

inducing the quotient morphism $C \to C/\langle \sigma_2 \rangle$; this establishes the isomorphism $C/\langle \sigma_2 \rangle \simeq E_{\sqrt{2}}$. Let $\alpha = 1 + \sqrt{2}$ so that

$$\sqrt{2}\, y^2 = x^3 - 35x^2 + 35x - 1 = (x-1)(x-\alpha^4)(x-\alpha^{-4})$$

is an affine equation for $E_{\sqrt{2}}$. Applying the Weil restriction functor to the morphism $C \to E_{\sqrt{2}}$ we obtain the morphism $\mathrm{WR}(C) \to \mathrm{WR}(E_{\sqrt{2}})$ defined over $\mathbb{Q}$. Composing with the inclusion of the diagonal $C \to \mathrm{WR}(C)$ we obtain a morphism $C \to \mathrm{WR}(E_{\sqrt{2}})$, inducing an isogeny $JC \to \mathrm{WR}(E_{\sqrt{2}})$, whose dual is ψ after extension of scalars from $\mathbb{Q}$ to $\mathbb{Q}(\sqrt{2})$.

By [Kuh88, Lemma, pag. 45] we conclude that the degree of the isogeny ψ is four (see also Remark 2.6 below). $\square$

Remark 2.5. In a similar way, the morphism induced by

$$(x,y) \mapsto \left(\left(\frac{x-\sqrt{-2}}{x+\sqrt{-2}} \right)^2, \frac{8\sqrt{-2}y}{3(x+\sqrt{-2})^3} \right)$$

determines isomorphisms

$$C/\langle \sigma_1 \circ \sigma_2 \rangle \simeq E'_{\sqrt{-2}} \qquad \text{and} \qquad C/\langle \sigma_1^3 \circ \sigma_2 \rangle \simeq E'_{-\sqrt{-2}}.$$

This implies the existence of a degree four isogeny $\mathrm{WR}(E'_{\sqrt{-2}}) \to JC$ defined over $\mathbb{Q}$.

Remark 2.6. Since the two elliptic curves $E_{\sqrt{2}}$ and $E_{-\sqrt{2}}$ are quadratic twists of each other, their two-torsion subgroups are canonically isomorphic; denote by T_2 the graph of this isomorphism in $E_{\sqrt{2}}[2] \oplus E_{-\sqrt{2}}[2]$. The group T_2 is the kernel of the isogeny ψ of Proposition 2.4 (this is an example of [Kuh88, Lemma, pag. 45]). In our example the two-torsion points on E (and hence also on $E_{\sqrt{2}}$ and on $E_{-\sqrt{2}}$) are

$$0 = (1, 0), \quad p_1 = (\alpha^4, 0), \quad p_2 = (\alpha^{-4}, 0), \quad p_3 = (\infty, 0),$$

and we have

$$\ker(\psi) = \{(0, 0), (p_1, p_2), (p_2, p_1), (p_3, p_3)\}.$$

We denote by E_1, E_2, R the images via ψ of $E_{\sqrt{2}} \times \{0\}$, $\{0\} \times E_{-\sqrt{2}}$ and the graph Δ of an isomorphism between the two twists.

Lemma 2.7. *With the previous notation, the following hold:*

(i) $\psi_{|E_{\sqrt{2}} \times \{0\}}$ *has degree* 1 *and* $\hat{\psi}_{|E_1}$ *has degree* 4 *(similarly for* $\{0\} \times E_{-\sqrt{2}}$ *and* E_2*);*

(ii) $\hat{\psi}_{|C}$ *has degree* 1*;*

(iii) $\psi_{|\Delta}$ *and* $\hat{\psi}_{|R}$ *have degree two.*

Proof. The composition of ψ with its dual $\hat{\psi}\colon JC \to \mathrm{WR}(E_{\sqrt{2}})$ is known to be the multiplication by 2. Thus the restriction of $\hat{\psi} \circ \psi$ to $E_{\sqrt{2}} \times \{0\}$, $\{0\} \times E_{-\sqrt{2}}$ and Δ has degree four.

By the description of $\ker(\psi)$ given in Remark 2.6, the preimage of E_1 by ψ is the union of $E_{\sqrt{2}} \times \{0\}$ and its translates by 2-torsion points. This gives (i).

Since $\hat{\psi}$ is a non-ramified covering, its restriction to C is clearly injective by Riemann-Hurwitz formula. This gives (ii).

By Remark 2.6 the kernel of ψ intersect the diagonal at two points, thus $\psi_{|\Delta}$ has degree two. This gives (iii). □

For the remainder of this section, we work over an algebraic closure of the ground field, and, to simplify the notation, we no longer distinguish between the elliptic curve E and its quadratic twists $E_{\sqrt{2}}$ and $E_{-\sqrt{2}}$.

Remark 2.8. The curve E appearing in Proposition 2.4 can be seen inside JC as follows. The hyperplane of equation:

$$\sqrt{2}(x_2 - x_4) + (x_1 + x_5) = 0$$

intersects X along the union of two conics C_1, C_2 and a smooth genus 1 curve. Each such conic intersects exactly four disjoint lines $R_1, \dots, R_4$ of X. Once these lines are contracted by the morphism mapping X to the Kummer quartic surface $\mathrm{Km}(JC)$, the image C_i' of each C_i passes through four nodes of the Kummer surface. The double cover

$$JC \to \mathrm{Km}(JC)$$

induces a double cover of C_i' branched along the four nodes. This is isomorphic to our elliptic curve E.

Before stating the next result, we recall some basic facts of lattice theory. Given a lattice S, its discriminant group is the finite abelian group $A_S := \mathrm{Hom}(S, \mathbb{Z})/S$ whose order equals $|\det(S)|$. If S is even, i.e. the quadratic form on S only takes even values, then there is an induced quadratic form $q_S\colon A_S \to \mathbb{Q}/2\mathbb{Z}$, called discriminant quadratic form.

If S is primitively embedded in a unimodular lattice L and $S^\perp \subset L$ is its orthogonal complement, then there is an isomorphism $A_S \cong A_{S^\perp}$. Moreover, up to this identification, the discriminant quadratic form on $A_{S^\perp}$ equals $-q_S$.

In what follows, U will denote the unique hyperbolic unimodular lattice of rank two and $U(m)$ the lattice obtained from U by multiplying its intersection matrix by m. Finally, E_8 denotes the negative definite root lattice associated to the Dynkin diagram of the corresponding type.

Theorem 2.9. *The Néron-Severi lattice and the transcendental lattice of the Jacobian surface JC are isomorphic to $U(2) \oplus (-2)$ and $U(2) \oplus (2)$ respectively.*

Proof. Since the j-invariant of E is not a rational integer, then E does not admit complex multiplication [Sil94, Corollary II.6.3.1]. Hence, by [KK01, Section 1] the transcendental lattice of the Kummer surface of $E \times E$ is isomorphic to $U(2) \oplus (4)$. Hence the transcendental lattice of $E \times E$ is isomorphic to

$$T_{E\times E} \cong U \oplus (2).$$

Now we will construct a basis of the Nerón-Severi lattice of JC. Let E_1, E_2, R be the images of $E \times \{0\}, \{0\} \times E$ and the diagonal in JC. In what follows we will indentify them with their classes in $\mathrm{NS}(JC)$. By Lemma 2.7 we have the following intersection numbers:

$$E_1 \cdot E_2 = 4, \quad R \cdot E_1 = R \cdot E_2 = 2, \quad C \cdot E_1 = C \cdot E_2 = 2.$$

In particular we deduce that C is numerically equivalent to $\frac{1}{2}(E_1 + E_2)$. The lattice N generated by E_1, R and $R + \frac{1}{2}(E_1 - E_2)$ has intersection matrix

$$N = U(2) \oplus (-2).$$

Observe that, since $\det(N) = 2^3$, the index of N in $\mathrm{NS}(JC)$ is either 1 or 2. We will prove that we are in the first case. The discriminant group A_N is isomorphic to $(\mathbb{Z}/2\mathbb{Z})^3$ and it is generated by the classes of $\frac{1}{2}E_1$, $\frac{1}{2}R$ and $\frac{1}{2}R + \frac{1}{4}(E_1 - E_2)$. Its discriminant quadratic form is

$$q_N = \begin{pmatrix} 0 & 1/2 \\ 1/2 & 0 \end{pmatrix} \oplus (-1/2).$$

If $D \in N$ is such that the class of $\frac{1}{2}D$ is in $\mathrm{NS}(JC)$, then $[\frac{1}{2}D] \in A_N$ is an isotropic vector for q_N. An easy check shows that the only non-zero isotropic vectors in A_N are the classes of $\frac{1}{2}E_1$ and $\frac{1}{2}R$. Since E_1 and R are integral curves with self-intersection zero, their classes are primitive in $\mathrm{NS}(JC)$. Thus we conclude that $[\frac{1}{2}D] = [0]$, that is $\frac{1}{2}D \in N$. This proves that $N = \mathrm{NS}(JC)$.

The transcendental lattice $T(JC)$ of JC is the orthogonal complement of N in the unimodular lattice $H^2(JC, \mathbb{Z}) \cong U^3$, so that it is a rank 3 lattice of signature $(2, 1)$ with $A_{T(JC)} \cong A_N$ and $q_{T(JC)} = -q_N$. By [Nik79, Theorem 1.13.2] there is a unique lattice, up to isomorphism, with these properties and one such lattice is $U(2) \oplus (2)$. This gives the statement. □

3. Two elliptic fibrations

By the results of the previous section we know that the surface X, which is the minimal resolution of the Kummer surface $\mathrm{Km}(JC)$, has Picard lattice of rank 19. There are two elliptic fibrations on X defined by:

$$\pi_{\pm}\colon X \to \mathbb{P}^1 \qquad (x_0 : \cdots : x_5) \mapsto (x_2 \pm x_4 : x_1 + x_5).$$

Observe that $\mathrm{Aut}(X)$ preserves the set of such fibrations. For example, $(x_2 - x_4 : x_1 + x_5)$ and $(x_2 + x_4 : x_1 - x_5)$ give fibrations defined by the same elliptic pencil on X due to the following relation in $I(X)$:

$$(x_1 + x_5)(x_1 - x_5) = 2(x_2 + x_4)(x_2 - x_4).$$

The types of the singular fibers of two fibrations π_+ and π_- coincide (since π_+ and π_- are exchanged by an automorphism of the surface) and they are

- 4 fibers of type I_4, whose components are the trivial lines of X,
- 2 fibers of type I_2, whose components are conics defined over $\mathbb{Q}(\sqrt{2})$, and
- 2 fibers of type I_2 whose components are conics defined over $\mathbb{Q}(\sqrt{-2})$.

The four squares which are fibers of π_+ are sections of π_- and viceversa. In particular, by [Shi00, Theorem 2.1] the torsion part of the Mordell-Weil group of both fibrations is isomorphic to $\mathbb{Z}/2\mathbb{Z} \oplus \mathbb{Z}/4\mathbb{Z}$. Thus, by our previous calculation of the Picard number of X, we have

$$\mathrm{MW}(\pi_{\pm}) \cong \mathbb{Z} \oplus \mathbb{Z}/2\mathbb{Z} \oplus \mathbb{Z}/4\mathbb{Z}.$$

We are ready to prove the following.

Theorem 3.1. *The set of rational points $X(\mathbb{Q})$ is Zariski-dense in X.*

Proof. The fibration π_- admits 16 sections coming from the irreducible components of the I_4-fibers of π_+; each of these sections is a line on X.

Let O be one of the 16 lines which are sections of π_-. Since the torsion part of the Mordell-Weil group of π_- has cardinality 8, there exists a line Q, which is a section of π_-, such that Q has infinite order with respect to the origin O. Since both the fibration π_- and the sections O, Q are defined over the rationals, we deduce that the multiples nQ are rational curves on X defined over $\mathbb{Q}$. Each such curve contains infinitely many rational points of X, and the theorem follows. □

Remark 3.2. As a consequence of Theorem 3.1 the subgroup $\mathrm{Aut}(X)_{\mathbb{Q}}$, of automorphisms of X defined over the rationals, is infinite. Indeed such a group contains a translation of the elliptic curve $X(\mathbb{Q}(t))$ of infinite order.

Observe that *any* rational point of X lies on a fiber of $\pi_{\pm}$ since both morphisms are defined over the rationals. The knowledge of the Mordell-Weil group of one of the two fibrations, let us say π_-, allows us to determine a basis for the Picard lattice of X.

Theorem 3.3. *The transcendental lattice of X is isomorphic to $U(4) \oplus (4)$, while the Picard lattice of X is isomorphic to and to $U(4) \oplus (-4) \oplus E_8^2$.*

Proof. By [Mor84, Proposition 4.3] the intersection matrix of the transcendental lattice $T(X)$ of X is obtained by multiplying by 2 the corresponding matrix of $T(JC)$. Hence $T(X)$ is isomorphic to $U(4) \oplus (4)$, by Theorem 2.9. In particular the discriminant group of $T(X)$ is isomorphic to $(\mathbb{Z}/4\mathbb{Z})^3$ with quadratic form given by:

$$q := \begin{pmatrix} 0 & 1/4 \\ 1/4 & 0 \end{pmatrix} \oplus (1/4).$$

The Picard lattice has signature $(1, 18)$ and discriminant group isomorphic to that of $T(X)$ with opposite quadratic form. By [Nik79, Theorem 1.13.2] a lattice with the previous properties is unique up to isomorphism, hence it is isomorphic to the lattice $U(4) \oplus (-4) \oplus E_8^2$ which has the same invariants. □

A computer calculation shows that the intersection matrix of all the lines and conics of X has rank 9 and discriminant group of order 2^6. Thus the Picard lattice is generated by the classes of such curves. Indeed a refined analysis shows that the classes of 17 lines plus the classes of two conics contained in the same fibration, the first defined on $\mathbb{Q}(\sqrt{2})$ and the second defined on $\mathbb{Q}(\sqrt{-2})$, still generate the Picard lattice.

Remark 3.4. Let μ_8 be the group of 8-th roots of unity. The Galois group G of the extension $\mathbb{Q}(\mu_8)/\mathbb{Q}$ acts on X and on its Picard lattice. Let C_1 and C_2 be two conics defined over $\mathbb{Q}(\sqrt{2})$ and $\mathbb{Q}(\sqrt{-2})$ respectively. Recall that the Picard lattice is generated by the classes of the lines together with those of C_1 and C_2. Since the action of G is trivial on the classes of the lines, we get a decomposition of G-modules

$$\mathrm{Pic}(X) \cong \mathrm{Pic}(X)_{\mathbb{Q}} \oplus L,$$

where the first factor has rank 17 and is spanned by the classes of all lines. The G-module L has rank 2 and is generated by the classes of C_1 and C_2. Observe also that since the two fibrations $\pi_\pm$ are defined over the rationals and each Mordell-Weil group is generated by classes of trivial lines, then any section of each fibration is in $\mathrm{Pic}(X)_{\mathbb{Q}}$.

Remark 3.5. A basis of the Picard group of X consists of classes of curves of the 19 elements set $S = \{L_1, \dots, L_{12}, C_1, C_2, C_3, R_1, \dots, R_4\}$ which satisfy the following properties. The L_i are lines contracted by π_- whose classes form an intersection lattice of type $A_3^{\oplus 4}$. Each line L_{3k} intersects two of the L_i lines. The C_i are disjoint conics contracted by π_-, while the R_i are lines which are sections of π_-. Consider the following sets: $S_1 = \{R_2, L_1, L_4, L_7, C_2, C_3\}$, $S_2 = \{R_2, L_6, L_{10}, C_1\}$, $S_3 = \{L_6, L_9, L_{11}, C_2, C_3\}$ and $S_4 = \{L_9, L_{10}, C_2\}$. Each line R_i intersects in one point each curve of the set S_i and is disjoint from any curve of $S \setminus S_i$. The intersection matrix of the classes of all such curves has rank 19 and Smith normal form with three non-zero entries equal to 4. Hence these curves form a basis of the Picard lattice by Theorem 3.3. Moreover the Mordell-Weil group of the fibration π_- is generated by the classes of R_1, R_2, R_3 and R_4.

3.1. An explicit sequence of infinitely many rational points.

We can use the results of the previous section to explicitly determine an element of infinite order on the generic fiber X_η of π_-. By putting $t = (x_4 - x_2)/(x_1 + x_5)$ and substituting into the equations (0.2), we get that X_η is a quartic curve contained in the three dimensional linear subspace of $\mathbb{P}^5$ of equations:

$$(2t^2 - 1)x_2 + (2t^2 + 1)x_4 - 2t\, x_5 = 0 \qquad (2t^2 - 1)x_1 - 4t\, x_4 + (2t^2 + 1)x_5 = 0.$$

The equations of X_η inside this space are the following:

$$(4t^4 - 8t^2 + 1)x_4^2 + (4t^3 + 2t)x_4x_5 - (4t^4 - 2t^2 + 1)x_5^2 + (12t^4 - 12t^2 + 3)x_0^2 = 0$$
$$(4t^4 - 8t^2 + 1)x_3^2 + (8t^3 + 4t)x_4x_5 - (4t^4 + 4t^2 + 1)x_5^2 + (16t^4 - 8t^2 + 4)x_0^2 = 0.$$

With the same notation as in Remark 3.5 we chose an origin for the Mordell-Weil group of X_η to be the point O which corresponds to the line R_1. Its coordinates are:

$$O := (-2t - 2 : 2t + 1 : -2t : -2t + 1 : 2t - 2 : 1).$$

Now we take the point Q which corresponds to the line R_2. Since R_1 and R_2 intersect at one point, then by [Shi90] the point Q has infinite order with respect to the origin O. Moreover we know Q to be the generator of the free part of the Mordell-Weil group of X_η. Its coordinates are the following:

$$Q := (2t + 2 : -2t - 1 : 2t : 2t - 1 : -2t + 2 : 1).$$

By calculating $2Q$ we obtain a one-parameter family of non-trivial rational solutions of the Büchi problem:

$$\begin{aligned} x_0 &= 12t^4 + 5t^2 - 1 & x_3 &= 8t^5 - 10t^3 - 6t \\ x_1 &= 8t^5 + 8t^4 + 22t^3 - 2t^2 + 2t + 2 & x_4 &= 8t^5 - 4t^4 - 2t^3 - 11t^2 + 2t - 1 \\ x_2 &= -8t^5 - 4t^4 + 2t^3 - 11t^2 - 2t - 1 & x_5 &= -8t^5 + 8t^4 - 22t^3 - 2t^2 - 2t + 2. \end{aligned}$$

To conclude this section we show that each fibrationπ_- and π_+ admits fibers defined over $\mathbb{Q}$ whose rank is > 1. Hence X contains more rational points that those which lie on the sections of both fibrations. In particular we prove the following.

Theorem 3.6. *There exists a Zariski-dense subset Ω of $X(\mathbb{Q})$ such that for any $p \in \Omega$ the fibers of π_+ and π_- through p have rank at least* 2.

Proof. For all but a finite number of points of $\mathbb{P}^1(\mathbb{Q})$, the corresponding fiber F of π_+ has positive rank by [Sil83] and gives a base change

$$\begin{tikzcd} Y \arrow[r] \arrow[d, "\varphi"'] & X \arrow[d, "\pi_-"] \\ F \arrow[r, "\pi_-|_F"'] & \mathbb{P}^1 \end{tikzcd}$$

whose new elliptic surface Y, defined over $\mathbb{Q}$, has Mordell-Weil group of rank at least 2, by [Sal09, Proposition 2.1]. In particular for all but a finite number of rational points of F, the fiber of φ has rank at least 2. Denote by $q \in F(\mathbb{Q})$ one of these points, and let $F_q := \varphi^{-1}(q)$ be the fiber of φ over q. Reasoning as before on the base change

$$\begin{tikzcd} Y_q \arrow[r] \arrow[d, "\varphi_q"'] & X \arrow[d, "\pi_+"] \\ F_q \arrow[r, "\pi_+|_{F_q}"'] & \mathbb{P}^1 \end{tikzcd}$$

we deduce that, for all but a finite number of rational points of F_q, the rank of the corresponding fiber of φ_q is at least 2. If $p \in F_q(\mathbb{Q})$ is such a point, then both the fibers of π_- and π_+ through p have rank at least 2. Varying $q \in F(\mathbb{Q})$ and $p \in F_q(\mathbb{Q})$ we see that the set of such points p is Zariski-dense in X. □

4. A MODULAR INTERPRETATION OF THE RATIONAL POINTS OF X

In this section we show that the surface X is a moduli space of rank two vector bundles on itself, giving a natural modular interpretation to the rational points on X. We do not know if there is a similar interpretation for the integral points on X.

Mukai studied the moduli spaces of stable vector bundles on abelian and K3 surfaces in its influential paper [Muk84]. In particular, he showed in [Muk84, Example 0.9] the existence of a beautiful correspondence between the K3 surfaces that are intersections of three quadrics in $\mathbb{P}^5$ and certain moduli spaces of vector bundles on the K3 surfaces themselves. The correspondence is obtained as follows. Let Y be a smooth complete intersection of three quadrics in $\mathbb{P}^5$; denote by $\mathbb{P}^2$ the net of quadrics containing X. In the net of quadrics $\mathbb{P}^2$, the discriminant locus Δ_Y is the locus of quadrics of rank at most 5: the discriminant Δ_Y is a plane curve of degree six. Denote by M_Y minimal resolution of the double cover of $\mathbb{P}^2$ branched over the sextic Δ_Y. The general point of M_Y corresponds to a quadric Q containing Y, together with the choice of a ruling by 2-planes of the quadric Q. Identifying smooth quadrics in $\mathbb{P}^5$ with the Grassmannian $Gr(2,4)$, we see that the two rulings by 2-planes of a quadric Q correspond to the second Chern classes of the rank 2 vector bundles on Q determined by the dual of the tautological vector bundle and by the universal quotient bundle. Thus, to each point of M_Y corresponding to a smooth quadric Q containing X together with a choice of ruling on Q, we associate a rank two vector bundle on X by restricting to X itself the rank two vector bundle on Q determined by the chosen ruling. This correspondence induces an isomorphism between M_Y and the irreducible component of the moduli space $\mathcal{M}(2,\mathcal{O}_X(1),2)$ of stable rank two vector bundles E on X with Chern classes $c_1(E) = [\mathcal{O}_X(1)]$ and $\deg(c_2(E)) = 4$.

Theorem 4.1. *The Büchi K3 surface X is isomorphic to the moduli space of stable rank two vector bundles on X itself with Chern classes $c_1(E) = [\mathcal{O}_X(1)]$ and $\deg(c_2(E)) = 4$.*

Proof. By the discussion above, it suffices to show that the double cover of $\mathbb{P}^2$ branched over the discriminant is isomorphic to X. First of all we give a birational map $f : X \to S \subseteq \mathbb{P}^3$, where S is a singular Kummer surface. Then we will show that the branch locus Δ_S of the projection $S \to \mathbb{P}^2$ from a node of S is projectively equivalent to the subscheme Δ of the singular quadrics in the net of quadrics defining X.

Let R be any line of X, for example that of equations:

$$x_1 + x_5 + 4x_0 \;=\; x_2 + x_5 + 3x_0 \;=\; x_3 + x_5 + 2x_0 \;=\; x_4 + x_5 + x_0 \;=\; 0.$$

Recall that there are exactly six lines $L_1, \dots, L_6$ of X incident to R. The birational map $f : X \to \mathbb{P}^3$ is defined by the linear system of quadrics of $\mathbb{P}^5$ with base locus $2R + \sum_{i=1}^{6} L_i$. In coordinates, if we denote by $z_0, \dots, z_3$ the four linear polynomials which define R, then f is given by:

$$\begin{aligned}
&6(z_0z_2 - z_1z_2 - z_0z_3 + z_2z_3)\\
&z_0z_1 - 2z_0z_2 + 3z_0z_3 - 4z_1z_3 + 2z_2z_3\\
&z_0z_1 - 2z_1z_2 - 3z_0z_3 + 4z_1z_3\\
&2z_0z_1 - 6z_0z_2 + 6z_1z_2 - 8z_1z_3 + 6z_2z_3.
\end{aligned}$$

The surface $S = f(X)$ has equation:

$$x_0^4 + 36x_1^4 + 36x_2^4 + x_3^4 - 4(x_0^2x_1^2 + x_0^2x_2^2 + x_1^2x_3^2 + x_2^2x_3^2) - 56x_1^2x_2^2 - \frac{14}{9}x_0^2x_3^2 = 0.$$

The sixteen nodes of S are defined over the rationals. The branch locus Δ_S of the projection $\pi\colon S \to \mathbb{P}^2 = V(x_3)$ from the node $(-3 : -1 : 0 : 3)$ is the union of the six tangent lines to the conic

$$9x_0^2 - 12x_0x_1 + 36x_1^2 - 32x_2^2 = 0$$

at the points $(-6 : 7 : 9)$, $(6 : -7 : 9)$, $(-6 : -1 : 3)$, $(6 : 1 : 3)$, $(-2 : -3 : 3)$, $(2 : 3 : 3)$. The scheme Δ of singular quadrics of the net which defines X is the union of the the six tangent lines to the conic:

$$x_0^2 + 8x_0x_1 - 34x_0x_2 + 16x_1^2 + 8x_1x_2 + x_2^2 = 0$$

at the points $(0 : -1 : 4)$, $(9 : 2 : 1)$, $(1 : -2 : 1)$, $(1 : 1 : 1)$, $(-4 : 1 : 0)$, $(1 : 2 : 9)$. An easy calculation shows that the unique projectivity $\sigma : \mathbb{P}^2 \to \mathbb{P}^2$ that maps the first four points of the first sextuple into the first four points of the second sextuple, maps the first set of six points into the second set of six points. Hence $\sigma(\Delta_S) = \Delta$, so that the corresponding double covers are isomorphic. □

Remark 4.2. In the proof of Theorem 4.1 we explicitly computed the discriminant of the net of quadrics and checked that the resulting K3 surface was isomorphic to the surface X. It is classically known that the K3 surfaces that are complete intersections of a net of quadrics and the double cover of the net of quadrics branched over the discriminant locus are isomorphic as soon as the surface contains a line; thus in our case we could have avoided the explicit computation, because the surface X contains 32 lines. A modern reference for this result is [MN04].

Over the field $\mathbb{Q}(\sqrt{-6})$ we can absorb more of the coefficients of the equation of the Kummer surface S in $\mathbb{P}^3$ to obtain the equation

$$9(x_0^4 + x_1^4 + x_2^4 + x_3^4) + 6(x_0^2x_1^2 + x_2^2x_3^2) + 6(x_0^2x_2^2 + x_1^2x_3^2) - 14(x_0^2x_3^2 + x_1^2x_2^2) = 0.$$

5. SUPERSINGULAR PRIMES FOR X

In this section we will prove a congruence property for the supersingular primes for the elliptic curve E of equation $y^2 = x^3 - 8x^2 - 2x$, isomorphic over the rational numbers to the elliptic curve of equation $y^2 = x^3 - 35x^2 + 35x - 1$.

Lemma 5.1. *The number field K_4 obtained by extending $\mathbb{Q}$ with the abscissas of all the 4-torsion points of the curve E is the 24th cyclotomic field $\mathbb{Q}(\mu_{24})$.*

Proof. The field K_4 is the splitting field of the polynomial

$$x(x^2 - 8x - 2)(x^2 + 2)(x^4 - 16x^3 - 12x^2 + 32x + 4).$$

The roots of the degree four factor are

$$4 + 2\sqrt{3} \pm (2\sqrt{6} + 3\sqrt{2}) \qquad \text{and} \qquad 4 - 2\sqrt{3} \pm (2\sqrt{6} - 3\sqrt{2}),$$

hence an easy calculation shows $\mathbb{Q}(\mu_{24}) = K_4$, as required. □

A calculation using Magma shows that the abelianization of the Galois group of the field $\mathbb{Q}(E[8])$ is the 24th cyclotomic field, but we do not need this more precise information: for us it is enough to know that the field $\mathbb{Q}(\mu_{24})$ is contained in $\mathbb{Q}(E[8])$.

In the next theorem we adapt a strategy that we learned from [Bra09]. In the proof we work with the 8-torsion subgroup, since the information coming from the 4-torsion elements is not enough for our purposes.

Theorem 5.2. *Let p be a supersingular prime for the elliptic curve E. Then $p \equiv 5$ or $23 \mod 24$.*

Proof. We concentrate on the Galois representation

$$\rho\colon \operatorname{Gal}(\bar{\mathbb{Q}}/\mathbb{Q}) \to \mathrm{GL}_2(\mathbb{Z}/8\mathbb{Z})$$

obtained by acting with the absolute Galois group on the set of 8-torsion points of the curve E. By Lemma 5.1 the image $G(8)$ of ρ admits a quotient $A(8)$ isomorphic to the Galois group of the extension $\mathbb{Q}(\mu_{24})/\mathbb{Q}$. Given a prime of good reduction p, the Frobenius automorphism acting on the reduction of X modulo p lifts to an element $\Phi_p \in \operatorname{Gal}(\bar{\mathbb{Q}}/\mathbb{Q})$. The image of Φ_p in the quotient $A(8) \cong (\mathbb{Z}/24\mathbb{Z})^*$ of the representation ρ is a congruence class modulo 24. If p is a supersingular prime for E, then Φ_p has null trace. We will show that the image in $A(8)$ of elements with null trace is congruent to either 5 or 23 modulo 24, concluding the proof. To prove the last step, we made use of Magma to show that the image $G(8)$ of the Galois representation ρ is generated by the following matrices:

$$\begin{pmatrix} 4 & 3 \\ 1 & 4 \end{pmatrix} \qquad \begin{pmatrix} 2 & 1 \\ 1 & 2 \end{pmatrix} \qquad \begin{pmatrix} 3 & 6 \\ 2 & 7 \end{pmatrix} \qquad \begin{pmatrix} 1 & 2 \\ 6 & 3 \end{pmatrix}.$$

The group $G(8)$ is a solvable group of order 2^6 and index $2^3 \cdot 3$ in $\mathrm{GL}_2(\mathbb{Z}/8\mathbb{Z})$. It contains 16 elements of null trace whose images in the quotient $A(8)$ form a subset S of cardinality 2. Since the primes 5 and 167 are supersingular for E and the second is congruent to 23 modulo 24, the images of Φ_5 and Φ_{167} in $A(8)$ exhaust the whole S. This concludes the proof. □

Theorem 5.3. *Let p be a supersingular prime for the elliptic curve E such that $p \equiv 23 \mod 24$. Then the number of points of the reduction of X modulo p equals $p^2 + 18p + 1$.*

Proof. Let JC be the Jacobian of C and let $Y := JC/\langle\sigma\rangle \subset \mathbb{P}^3$ be its quotient with respect to the involution $\sigma\colon w \mapsto -w$. Observe that $\#X(\mathbb{F}_p) = \#Y(\mathbb{F}_p) + 16p$ for any prime $p \geq 5$ since all the sixteen singular points of Y are defined over $\mathbb{Z}$. Consider two subsets of JC over $\mathbb{F}_{p^2}$:

$$S_1 := \{w \in JC(\mathbb{F}_{p^2}) : \Phi_p(w) = w\} \qquad S_2 := \{w \in JC(\mathbb{F}_{p^2}) : \Phi_p(w) = -w\},$$

where Φ_p is the Frobenius map. We have $\#Y(\mathbb{F}_p) = \frac{1}{2}(\#S_1 + \#S_2)$. We now want to show that $S_1 = JC(\mathbb{F}_p)$ and that

$$\#S_2 = \#J\tilde{C}(\mathbb{F}_p), \tag{5.1}$$

where $\tilde{C}$ is the quadratic twist of C. The first equality is obvious. About the second equality, let $y^2 = f(x)$ be an equation for C and $dy^2 = f(x)$ be an equation for $\tilde{C}$. Consider the isomorphism

$$f\colon C \to \tilde{C} \qquad (x, y) \mapsto \left(x, \frac{1}{\sqrt{d}}y\right)$$

defined over $\mathbb{F}_{p^2}$ and observe that $\Phi_p \circ f = \tilde{\iota} \circ f \circ \Phi_p$, where $\tilde{\iota}$ is the hyperelliptic involution of $\tilde{C}$. Let $f_*\colon JC \to J\tilde{C}$ be the push-forward map defined by $f_*([\sum n_i q_i]) = [\sum n_i f(q_i)]$. Then by the above equality we get $\Phi_p \circ f_* = -f_* \circ \Phi_p$.

To prove (5.1) it is enough to show that $f_*(S_2) = J\tilde{C}(\mathbb{F}_p)$, which immediately follows from

$$\Phi_p(f_*(w)) = -f_*(\Phi_p(w)) = f_*(w),$$

where $w \in S_2$.

Assume now that p is a supersingular prime for E such that $p \equiv 23 \pmod{24}$. Hence the Jacobian variety JC is isogenous to $E_{\sqrt{2}} \times E_{-\sqrt{2}}$, by Proposition 2.4 and the fact that 2 is a square in $\mathbb{F}_p$. This gives

$$\#S_1 = \#JC(\mathbb{F}_p) = (p+1)^2.$$

Moreover the trace of the Frobenius morphism Φ_q, with $q = p^n$, on $H^1_{\text{ét}}(JC, \mathbb{Q}_l)$ equals the trace of the Frobenius morphism on the first étale cohomology group of $E_{\sqrt{2}} \times E_{-\sqrt{2}}$. Since $E_{\pm\sqrt{2}} \mod p$ is supersingular then its Zeta function is:

$$Z(E_{\pm\sqrt{2}}, t) = \frac{qt^2+1}{(1-t)(1-qt)}.$$

By means of the isogeny $JC \sim E_{\sqrt{2}} \times E_{-\sqrt{2}}$ we deduce that the Zeta Function of C is

$$Z(C,t) = \frac{(qt^2+1)^2}{(1-t)(1-qt)}. \tag{5.2}$$

Observe that $\#C(\mathbb{F}_p) = \#\tilde{C}(\mathbb{F}_p)$ since $\#C(\mathbb{F}_p) + \#\tilde{C}(\mathbb{F}_p) = 2p+2$ and $\#C(\mathbb{F}_p) = p+1$, by the above calculation of the Zeta function of C. Hence C and $\tilde{C}$ have the same Zeta function, since the two curves have the same number of points over $\mathbb{F}_p$, and same number of points over $\mathbb{F}_{p^2}$, being isomorphic over this field. This gives $\#JC(\mathbb{F}_p) = \#J\tilde{C}(\mathbb{F}_p) = (p+1)^2$, so that $\#Y(\mathbb{F}_p) = (p+1)^2$ and $\#X(\mathbb{F}_p) = (p+1)^2 + 16p$ as claimed. □

Remark 5.4. Theorem 5.2 shows that the supersingular primes for the curve E are congruent to 5 or 23 modulo 24. For those congruent to 5 modulo 24 the Zeta functions of C and of $\tilde{C}$ coincide and are equal to

$$Z(C,t) = Z(\tilde{C},t) = \frac{(qt^2-1)^2}{(1-t)(1-qt)},$$

analogous to formula (5.2). We verified this formula for the primes 5, 149, 173, 461, 1229, 2213, 2237 that are the first 7 supersingular primes of E congruent to 5 modulo 24. As a consequence, the number of points of X modulo a prime p congruent to 5 modulo 24 and supersingular for E is $p^2 + 14p + 1$.

References

[Bea96] Arnaud Beauville, *Complex algebraic surfaces*, Second, London Mathematical Society Student Texts, vol. 34, Cambridge University Press, Cambridge, 1996. ↑5

[BLR90] Siegfried Bosch, Werner Lütkebohmert, and Michel Raynaud, *Néron models*, Ergebnisse der Mathematik und ihrer Grenzgebiete (3) [Results in Mathematics and Related Areas (3)], vol. 21, Springer-Verlag, Berlin, 1990. ↑6

[BCP97] Wieb Bosma, John Cannon, and Catherine Playoust, *The Magma algebra system. I. The user language*, J. Symbolic Comput. **24** (1997), no. 3-4, 235–265. Computational algebra and number theory (London, 1993). ↑5

[Bra09] Julio Brau, *Congruence conditions on supersingular primes*, Master's thesis, Mathematisch Instituut, Universiteit Leiden, 2009. ↑14

[KK01] Jonghae Keum and Shigeyuki Kondō, *The automorphism groups of Kummer surfaces associated with the product of two elliptic curves*, Trans. Amer. Math. Soc. **353** (2001), no. 4, 1469–1487 (electronic). ↑8

[Kuh88] Robert M. Kuhn, *Curves of genus* 2 *with split Jacobian*, Trans. Amer. Math. Soc. **307** (1988), no. 1, 41–49. ↑6, 7

[MN04] Carlo Madonna and Viacheslav V. Nikulin, *On a classical correspondence between $K3$ surfaces. II*, Strings and geometry, 2004, pp. 285–300. ↑13

[Mor84] D. R. Morrison, *On $K3$ surfaces with large Picard number*, Invent. Math. **75** (1984), no. 1, 105–121. ↑9

[Muk84] Shigeru Mukai, *Symplectic structure of the moduli space of sheaves on an abelian or $K3$ surface*, Invent. Math. **77** (1984), no. 1, 101–116. ↑12

[Nik75] V. V. Nikulin, *Kummer surfaces*, Izv. Akad. Nauk SSSR Ser. Mat. **39** (1975), no. 2, 278–293, 471. ↑5

[Nik79] ______, *Integer symmetric bilinear forms and some of their geometric applications*, Izv. Akad. Nauk SSSR Ser. Mat. **43** (1979), no. 1, 111–177, 238. ↑8, 10

[PPV10] H. Pasten, T. Pheidas, and X. Vidaux, *A survey on Büchi's problem: new presentations and open problems*, Zap. Nauchn. Sem. S.-Peterburg. Otdel. Mat. Inst. Steklov. (POMI) **377** (2010), no. Issledovaniya po Teorii Chisel. 10, 111–140, 243. ↑1, 2

[Sal09] Cecília Salgado, *Rank of elliptic surfaces and base change*, C. R. Math. Acad. Sci. Paris **347** (2009), no. 3-4, 129–132. ↑11

[Shi00] Ichiro Shimada, *On elliptic $K3$ surfaces*, Michigan Math. J. **47** (2000), no. 3, 423–446. ↑9

[Shi90] Tetsuji Shioda, *On the Mordell-Weil lattices*, Comment. Math. Univ. St. Paul. **39** (1990), no. 2, 211–240. ↑11

[Sil83] Joseph H. Silverman, *Heights and the specialization map for families of abelian varieties*, J. Reine Angew. Math. **342** (1983), 197–211. ↑11

[Sil94] ______, *Advanced topics in the arithmetic of elliptic curves*, Graduate Texts in Mathematics, vol. 151, Springer-Verlag, New York, 1994. ↑8

Departamento de Matemática, Universidad de Concepción, Casilla 160-C, Concepción, Chile
E-mail address: `michela.artebani@gmail.com, antonio.laface@gmail.com`

Mathematics Institute, Zeeman Building, University of Warwick, Coventry CV4 7AL, United Kingdom
E-mail address: `adomani@gmail.com`